

CROPS



BONSILAGE FORTE is developed specifically for all types of wet and hard-to-ensile forages in the lower dry matter ranges, which often exhibit high ash content with a high buffering capacity. **BONSILAGE FORTE** quickly lowers the pH, effectively inhibits clostridia growth and decreases the risk of butyric acid fermentation.

TYPE

Biological and water soluble silage additive

DOSAGE

At least 300,000 CFU/g fresh matter (FM) of forage

DRY MATTER RANGE OF CROPS

Grass haylage, clover grass haylage, alfalfa haylage, small grain silage: 22-35% DM

STRAINS

Lactobacillus plantarum,
Pediococcus acidilactici,
Lactococcus lactis

INGREDIENTS

Selected strains of homofermentative lactic acid bacteria, dextrose

ACTIVE SUBSTANCE


Lactic acid bacteria not less than 1.36×10^{11} CFU/g product

- » *Lb. plantarum* and *Pc. acidilactici* quickly lower the pH level by producing lactic acid in the front-end fermentation cycle.
- » *Lc. lactis* is known for its active clostridia inhibition to reduce the risk of butyric acid fermentation, thereby reducing proteolysis and thereby retaining more of the harvested forage protein.
- » **BONSILAGE FORTE** converts damp, raw material into hygienically nutrient-rich forages with excellent palatability.
- » **BONSILAGE FORTE** is specially developed for all types of wet and hard-to-ensile grass, alfalfa and small grain silages in the lower dry matter range.

RESEARCH

We conduct extensive on-farm research and feeding trials to ensure the highest level of performance from **BONSILAGE** products. This study shows how the lactic acid bacteria (LAB) contained in **BONSILAGE FORTE** quickly inhibited the level of active clostridia in grass silage (first cut, 26-32% DM) and then kept the level relatively low for 90 days.

Clostridia counts after various silo ripening times





SMALL CAN (200 G)
100 tons FM forage



LARGE CAN (1 KG)
500 tons FM forage

DIRECTIONS FOR USE

1. Fill a bucket with clean, cold (below 60 °F), unchlorinated water. Use at minimum 1 gallon of water per can.
2. Add the BONSILAGE product into the mixing bucket.
3. Dissolve the BONSILAGE product uniformly in the bucket.
4. Add more water to achieve desired application volume.



APPLICATION & OUTPUT

- » Apply 2 g of BONSILAGE FORTE equally to 1 ton of fresh matter (FM) forage, based on individual application rate and type of available applicator.
- » Avoid heating the solution during application try to stay below 70°F to preserve the LAB, and allow them the best possible performance.
- » Small can (200 g) will sufficiently treat 100 tons FM forage, large can (1 kg) will sufficiently treat 500 tons FM forage.
- » Do NOT add acids, salts or other substances, as they could reduce the number of viable bacteria in the product.

STORAGE OF PRODUCT

- » Store unopened bottles in a cool, dry place away from direct sunlight.
- » Use the entire bottle when opened.
- » The prepared solution can be stored for up to 48 hours if kept below 70°F.

BONSILAGE FORTE contains noble LAB strains that are preserved by the latest freeze-dried conservation technology. This allows all BONSILAGE products to be stored at room temperature, so freezer storage is NOT necessary. BONSILAGE FORTE comes in sealed plastic cans and has a 24-month shelf life from production date. Our sturdy packaging ensures high-quality protection against environmental influences and allows for convenient mixing with water.

FOR MORE INFORMATION:

888 580-7797
bonsilageusa.com
info@bonsilageusa.com

PLEASE NOTE:

BONSILAGE products are the most widely used silage inoculants in Europe. Our products contain living, specifically selected lactic acid bacteria (LAB) produced by Lactosan, which is a sister company to PROVITA SUPPLEMENTS and a leader in scientific selection and production of LAB for silage and probiotics in animal feed. Our access to such highly sought-after bacteria results in superior forage quality and feeding value.

BONSILAGE FORTE contains a balanced mix of highly active homofermentative lactic acid bacteria strains. With a well-managed ensiling process, accurate dosing and sufficient compaction of the forage, BONSILAGE FORTE can improve silage quality and reduce the risk of reheating. The target density for proper fermentation should be a minimum of 15 lbs DM/ft³. For complete fermentation, the silage should be stored a minimum of 3 weeks before start of feed out.